



Pepin

and the Hepatitis B *Vaccine*



TEXAS TECH
UNIVERSITY
HEALTH SCIENCES CENTER
AT EL PASO



Pepin

and the
Hepatitis B
Vaccine



Pepin you
are so big
today!


You are
one month
old today.

We are going
to see your
pediatrician.

Pepe?

Are you ok?





The hepatitis B virus can cause a short-term illness called acute that leads to:

Diarrhea and vomiting.



Loss of appetite.




Tiredness.




*Jaundice
(yellow skin
or eyes).*



*Pain in muscles,
joints, and stomach.*





But you will be protected by receiving the vaccine today.

Pepe, we need to get ready to go.

ZZZ



This disease can also cause long-term illness called chronic that leads to:



Liver damage (cirrhosis).



Liver cancer and




Death.






About one-and-a-quarter million people in the U.S. have chronic Hepatitis B Viral infection.




Each year approximately 80,000 people, mostly adults, become infected with the virus.




More than 11,000 people require hospitalization, and between 4,000 and 5,000 people die from chronic Hepatitis B.






Pepe do you know that the vaccine that Pepin will receive today...




...is in fact the first anti-cancer vaccine because it can prevent the form of liver cancer associated with this virus.





Quick, get ready,
we need to go and
protect Pepin.



I called my friend
Megan to go with me
because you were
not waking up.




We can all
go together.



I love to go
with you, Pepin...
...Pepe, and my
little cat.





I want to know more about the vaccine.

Hepatitis B virus is spread through contact with the blood and body fluids of an infected person.



A person can get infected by having unprotected sex with an infected person...


...or by sharing needles when injecting illegal drugs or...




...by being stuck with a blood contaminated needle.




In addition during birth when the virus passes from an infected mother to her baby.



One third of the people who are infected with Hepatitis B in the United States don't know how they got it.




Everyone under 18 years of age should get this vaccine, along with anyone 18 and older who is at risk.





Should I receive the vaccine?



Adults who
are at risk include
the following:




*People who have more
than one sex partner
in 6 months.*



*Men who have sex
with other men.*

*People who
have sex with an
infected person.*




*People who inject
illegal drugs.*

Health care and public safety workers who might be exposed to infected blood or body fluids.



Persons who share a household with someone who has chronic Hepatitis B.



Patients on hemodialysis.



I do not fit in any of these categories.














Pepin what is the matter?



You only need to get three doses of hepatitis B vaccine according to the following schedule.

If a dose is missed or behind schedule, there is not a need to start over instead it is indicated to get the next dose as soon as possible.

Hepatitis B Vaccination Schedule		Who?		
		Infants		Older child, adolescent, or adult
When?	First Dose	Mother infected with Hepatitis B Virus (HBV) 	Mother not infected with HBV 	
	Second Dose	Within 12 hours of birth 	Birth-2 months of age 	Anytime
	Third Dose	1-2 months of age 	1-4 months of age (at least 1 month after first dose) 	1-2 months after first dose
		6 months of age 	6-18 months of age 	4-6 months after first dose




First Dose

The first dose should be given:



Within 12 hours of birth to infants whose mothers are infected with Hepatitis B.



Anytime between birth and 2 months of age for infants whose mothers are not infected.


Anytime for older children, adolescents and adults.






Second Dose

Between 1 and 2 months of age to children of infected mothers with Hepatitis B.



Between 1 and 4 months of age, but at least 1 month after the first dose, for children of mothers who are not infected...

...and 1 to 2 months after the first dose for older children, adolescents and adults.




Third Dose

The third dose should be given:



At 6 months of age to children of infected mothers with Hepatitis B Virus.

At 6 to 18 months of age for children of mothers who are not infected, and...




...4 to 6 months after the first dose for older children, adolescents and adults.


Again, the second dose must be given at least 1 month after the first dose.



Finally the third dose must be given at least 2 months after the second dose and at least 4 months after the first.



The third dose should not be given to infants under 6 months of age...



...because this could reduce long-term protection.





Adolescents 11 to 15 years of age might need only two doses of Hepatitis B vaccine, separated by 4-6 months.



Ask your health care provider for details.












Have ever had a life-threatening allergic reaction to baker's yeast the kind used for making bread or...








...to a previous dose of the vaccine.









Wheezing.




Hives.



Paleness.



Weakness.



*Fast heart
beat or*




Dizziness.



*In the event
of a serious reaction,
call a doctor or get the person
to a doctor right away.*

*The possibility
of these reactions
are very rare.*





*Ask your doctor,
nurse, or health department to file a*

Vaccine Adverse Event Reporting System Form

call the VAERS yourself at

1-800-822-7967

or visit their website at

www.vaers.org



In the rare event
that you or your child has
a serious reaction to
a vaccine...

...a federal program has
been created to help you pay
for the care of those who
have been harmed.



National Vaccine Injury

Compensation Program

call

1-800-338-2382

or visit the program's website at

www.hrsa.gov/osp/vicp



Well let us go home.



Hmmm,
maybe Pepe isn't quite
ready yet.

Director:
Gilbert Handal, M.D.

Producer:
Marie A. Leiner, Ph.D.

Storywriter:
Gilbert Handal, M.D.

Adaptation:
Marie A. Leiner, Ph.D.
Miriam Silva, M.D.

Editor:
Carlos González H., Ph.D.

Coordinator:
Carmen Avila, B.A.

Graphic Design:
Leader Eleuterio Meza, B.A.
Fabián Luna, B.A.
Adrián Durán, B.A.

Pepin *and the* **Hepatitis B Vaccine**



The production of these resources was supported by Grant Number U21/CCU624220-01 from the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not represent the official views of the CDC.